MALE SYMPTOM MONITOR

Name:	Date:	
Occupation:	Age:	
Complaints:	1.	
	2.	
	3.	
SURGICAL HISTOR	<u>Y</u> :	
Abdominal: When:		
Pelvic: When:		
	MS: Please put an X next to the statements that be	
My incontinence is associat	ed with activities such as sneezing, running or coug	hing daily weekly
S My incontinence is precede U	d by a strong sensation that feels uncontrollable	daily weekly
-	ed with frequency of urination during the day (>5-7	' X/day) # times per day
My bladder troubles cause t	frequent nighttime urination	# times/night
= =	ed with frequent nighttime bedwetting	# times/week
My incontinence requires m	ne to wear pads	# pads/day
My bladder troubles include	e incomplete emptying	Yes No Sometimes
R I have pain when I urinate		Yes No Sometimes
I have to strain when I urina	ate	Yes No Sometimes
TP I have leakage during interc	ourse	Yes No Sometimes
I had problems with urination	on during my childhood	Yes No
Urinary Urgency without ur	ine loss	Yes No
Fluid Intake in 24 hours:		
# cups of soffoo/day	# support water/day # support to a /day	# support other fluids /day

Pelvic Health Solutions

Restoring Pelvic Health through Physiotherapy

BOWEL HISTORY:
Frequency: /week
Fecal Incontinence: Yes No Stool Consistency: Loose Soft/formed Hard Varies
Fecal Urgency: Yes No
Constipation: Yes No
MEDICAL HISTORY:
Urinary Tract Infections: Yes No Antibiotics Recently? Yes No
Smoking: Yes No #packs/day
Chronic Cough: Yes No
Do you get blood in your urine: Yes No
Allergies (including latex):
Height: ft In. Weight: Ibs BMI:(therapist)
Back Problems: Yes No If yes, please ask the receptionist for the Pelvic Girdle Pain Assessment
Neck Problems: Yes No Chronic? Yes No
Have you ever been treated for depression? Yes No
SEXUAL HISTORY:
Last PSA Score: When?
Last digital rectal exam?
Prostate Fluid expressed and tested? Yes No
Do you have painful erections? Yes No
Can you achieve a satisfactory erection? Yes No
Do you have premature ejaculation? Yes No
Do you have pain during intercourse? Yes No
On a scale from 1-10, please circle and rate your current pain/discomfort
1 2 3 4 5 6 7 8 9 10

DASS Questionnaire

(**Reference:** Loribond et al. The structure of negative emotional states; comparison of the DASS with the Beck Inventories. Beh. Res. Ther 1995; 33:335-342)

Please read each statement and circle a number, 0, 1, 2 or 3, which indicates how much the statement applied to <u>over the past week</u>. There are no right or wrong answers. Do not spend too much time on any statement.

- 0.....Did not apply to me at all
- 1.....Applied to me to some degree or some of the time
- 2.....Applied to me a considerable degree, or a good part of the time
- 3.....Applied to me very much, or most of the time

I found it hard to wind down	5	0	1	2	3
I was aware of dryness of my mouth	Α	0	1	2	3
I could not seem to experience any feeling at all	D	0	1	2	3
I experienced breathing difficulty (e.g. excessively rapid breathing, breathlessness in the absence of physical exertion)	Α	0	1	2	3
I found it difficult to work up the initiative to do things	D	0	1	2	3
I tended to over-react to situations	S	0	1	2	3
I experienced trembling (e.g. in the hands)	Α	0	1	2	3
I felt that I was using a lot of nervous energy	S	0	1	2	3
I was worried about situations in which I might panic and make a fool of myself	Α	0	1	2	3
I felt that I had nothing to look forward to	D	0	1	2	3
I found myself getting agitated	S	0	1	2	3
I found it difficult to relax	S	0	1	2	3
I felt down-hearted and blue	D	0	1	2	3
I was intolerant of anything that kept me from getting on with what I was doing	S	0	1	2	3
I felt I was close to panic	Α	0	1	2	3
I was unable to become enthusiastic about anything	D	0	1	2	3
I felt I was not worth much as a person	D	0	1	2	3
I felt that I was rather touchy	S	0	1	2	3
I was aware of the action of my heart in the absence of physical exertion (e.g. sense of heart rate increase, heart missing a beat)	Α	0	1	2	3
I felt scared without any good reason	Α	0	1	2	3
I felt that life was meaningless	D	0	1	2	3

a	A	T.
_	A =	I 1 —
S =	A -	$\mathbf{D} =$

Assessment Symptom Outcome Measure

The following questions are reproduced with permission from the International Pelvic Pain Society, <u>www.pelvicpain.org</u>

Please read each of the following statements and circle the number that best represents your symptoms:

0 = no pain 1 = mild pain 2 = mild-moderate pain 3 = moderate4 = moderate-severe 5 = severe symptoms Pain:

<u>Pain:</u>						
How would you rate your present pain	0	1	2	3	4	5
Pain when lifting	0	1	2	3	4	5
Pain when sitting	0	1	2	3	4	5
Pain when walking	0	1	2	3	4	5
Pain while doing physical activity	0	1	2	3	4	5
Deep pain with sexual intercourse or sexual activity	0	1	2	3	4	5
Pelvic pain lasting hours or days after sexual activity	0	1	2	3	4	5
Pain when bladder when full	0	1	2	3	4	5
Pain with urination	0	1	2	3	4	5
Muscle or joint pain	0	1	2	3	4	5
Abdominal pain	0	1	2	3	4	5
Backache	0	1	2	3	4	5
Pain when wearing tight clothing	0	1	2	3	4	5
Pain with bowel movement	0	1	2	3	4	5
Pain after bowel movement	0	1	2	3	4	5
A falling-out feeling or a feeling of pressure in the pelvis	0	1	2	3	4	5
Bladder Symptoms:		1	2	3	4	5
Loss of urine when coughing, sneezing, lifting or laughing	0	1	2	3	4	5
Frequency of urination versus the normal of once every 2-3 hours	0	1	2	3	4	5
Urgency or need to urinate with little warning	0	1	2	3	4	5
Loss of urine due to strong urge	0	1	2	3	4	5
Difficulty initiating urine stream	0	1	2	3	4	5
Urine stream stops and starts	0	1	2	3	4	5
Nighttime urinary frequency	0	1	2	3	4	5
Incomplete emptying of urine	0	1	2	3	4	5
Bowel Symptoms:	0	1	2	3	4	5
Constipation (fewer than 3 bowel movements/week)	0	1	2	3	4	5
Bowel frequency (more than 3 bowel movements/day)	0	1	2	3	4	5
Incomplete emptying of bowel	0	1	2	3	4	5
Urgency or need to have a bowel movement with little warning	0	1	2	3	4	5
Abdominal bloating or fullness	0	1	2	3	4	5
Lumpy or hard stool consistency	0	1	2	3	4	5
Needing to strain to achieve bowel movement	0	1	2	3	4	5
Fecal incontinence	0	1	2	3	4	5

Since your symptoms began, how much as your lifestyle been affected?

0 = no effect 1 = mild affect 2 = mild-moderate affect 3 = moderate affect 4 = moderate-severe 5 = substantive change

Effect on Daily Life:

Symptoms or pain limits or interferes with work or school	0	1	2	3	4	5
Symptoms or pain limits or interferes with social activities	0	1	2	3	4	5
Symptoms or pain limits or interferes with exercise routine	0	1	2	3	4	5
Symptoms or pain limits or interferes with lifting, cleaning, carrying, shopping, etc	0	1	2	3	4	5
Symptoms or pain limits or interferes with recreational and/or athletic activities	0	1	2	3	4	5
Symptoms or pain limits or interferes with sexual activity	0	1	2	3	4	5
Symptoms or pain limits or interferes with sleep	0	1	2	3	4	5
Symptoms or pain cause unexplained mood changes	0	1	2	3	4	5
Difficulty getting an erection (even when aroused)	0	1	2	3	4	5
Difficulty achieving orgasm (even when aroused)	0	1	2	3	4	5

Total: _____/ 220